

	Hospital Emergency Eye Service	CUES
<b>Trauma / Pain / Red Eye</b>	<ul style="list-style-type: none"> <li>• Chemical injury (Alkaline/Acid)</li> <li>• Penetrating injury</li> <li>• Lid laceration</li> <li>• Proptosis with visual loss / pain</li> <li>• Swollen lids (with pyrexia, +/- diplopia, distorted vision)</li> <li>• Red eye + LOV + Severe Pain +/- Nausea +/- photophobia</li> <li>• CL Wearer + Red painful eye</li> </ul>	<ul style="list-style-type: none"> <li>• Blunt trauma</li> <li>• Corneal foreign body (FB)</li> <li>• Corneal abrasion</li> <li>• Sub tarsal/conjunctival FB</li> <li>• New droopy lid/ ptosis</li> <li>• Pain on ocular movement</li> <li>• Swollen lids (normal vision, a pyrexia)</li> <li>• Red Eye, +/- Visual changes</li> <li>• Mild - moderate pain</li> <li>• Gritty eye</li> <li>• Discharge / watery eye</li> <li>• Itch</li> </ul>
<b>Vision</b>	<ul style="list-style-type: none"> <li>• Sudden loss of vision &lt;12 hours</li> <li>• New / sudden onset diplopia</li> <li>• New Flashes/floaters + Curtain Effect</li> </ul>	<ul style="list-style-type: none"> <li>• Sudden loss of vision &gt;12 hrs</li> <li>• Unilateral VF defect</li> <li>• New Flashes/floaters</li> <li>• Change in vision &lt;2/52</li> </ul>
<b>Headache / Neuro</b>	Headache + Painful scalp, Brow pain, Jaw Claudication, Painful temples (Pt. over 50 +/- diplopia or LOV)	<ul style="list-style-type: none"> <li>• New headache/migraine + ocular symptoms</li> <li>• New onset anisocoria with no other symptoms</li> <li>• Bell's palsy</li> </ul>
<i>This list is not exhaustive. Practitioners should always apply their clinical judgement when deciding on the appropriate clinical pathway</i>		
	Hospital Emergency Eye Service	CUES
<b>Condition</b>	<ul style="list-style-type: none"> <li>• Endophthalmitis</li> <li>• Papilloedema / swollen discs</li> <li>• Acute angle closure glaucoma</li> <li>• Suspected periocular / ocular malignancy</li> <li>• Compartmental Syndrome</li> <li>• Anterior Uveitis</li> </ul>	